

BULLETIN

OF THE

BRITISH ORNITHOLOGISTS' CLUB.

EDITED BY

N. B. KINNEAR.



VOLUME XLVI.
SESSION 1925-1926.

LONDON:

WITHERBY & CO., 326 HIGH HOLBORN.

1926.

Mr. GREGORY M. MATHEWS sent the following :—

Notochibia monica, new name for *Chibia densa* Sharpe, Cat. Birds Brit. Mus. vol. iii. p. 241 (1877). Timor (not Gray, 1870).

Meliphaga lombokia, new name for *Ptilotis virescens* Wallace (*cf.* Cat. Birds Brit. Mus. vol. ix. pl. vii. fig. 1), not Vieillot, 1817.

Scaephaethon rubricauda roseotincta, new name for *Phaëthon rubricauda erubescens* Rothschild 1900, not Gray 1844.

Poliolimnas cinereus collingwoodi, new name for *P. c. ocellaris* Ingram, 1911, preoccupied as a synonym of *cinereus* Vieillot.

Hæmatæna richmondæna, new name for *Columba melanocephala* Pennant 1769, not Brünnich 1764.

Nycticorax caledonicus pelewensis, subsp. nov.

Differs from *N. c. hilli* Mathews in being smaller and more dusky in tint, the inner webs of the quills being dull chestnut.

Distr. Pelew Islands.

Type in the British Museum. Pelew Islands. Cohen Coll. Brit. Mus. Reg. 78.10.29.18.

Poliolimnas cinereus moluccanus, subsp. nov.

Differs from *P. c. cinereus* (V.) in being lighter generally ; it is also smaller in bill and wing measurements.

Type in the Tring Museum. ♀ ad. Mt. Fogi, W. Buru. 18.ii.1902. Kuhn Coll. No. 4955.

Distr. Moluccas to Key Islands and Lesser Sunda Islands.

Pseudochlamydera, gen. nov.

Differs from *Chlamydera* Gould in lacking the nuchal crest.

Type, *Chlamydera lauterbachii* Reichenow.